

Beau lines associated with COVID-19

ÁNGEL ROLANDO GALVÁN-MARÍN • BOGOTÁ (COLOMBIA)

DOI: <https://doi.org/10.36104/amc.2021.2055>



Figure 1. Beau lines: Horizontal grooves (arrows) 4 mm from the proximal nail folds..

A 33-year-old male healthcare worker in an intensive care unit had horizontal grooves on all his nails. These grooves were 4 mm from the proximal nail fold (Figure 1). Four months ago, in the course of his work, he developed a fever, cough, diarrhea, arthralgia and hyposmia for eight days, but did not require hospitalization. He was ultimately diagnosed with COVID-19 through a positive nose swab PCR test. He had no traumas or other concurrent febrile disease.

Beau lines are caused by a temporary interruption of the growth of the proximal nail matrix (1). They generally occur after episodes of cardiac arrest, chemotherapeutic agents, febrile illnesses, trauma and Raynaud's disease (2). Since the beginning of the COVID-19 pandemic, reported dermatological findings have included maculopapular, vesicular and urticarial rashes (3). This image is an invitation to actively seek a new spectrum of COVID-19 manifestations.

Dr. Ángel Rolando Galván Marín: Especialista en Medicina Interna, Nefrología y Epidemiología. Nefrólogo Bbraun Avitum Renal Care. Nefrología Crítica Clínica Juan N. Corpas. Facultad de Medicina y Ciencias de la Salud Universidad del Rosario. Bogotá, D.C. (Colombia).

Correspondencia: Dr. Ángel Rolando Galván Marín. Bogotá (Colombia).

E-Mail: agalvanmed@hotmail.com

Received: 25/X/2020 Accepted: 19/V/2021

References

1. Jandial A, Mishra K, Prakash G, & Malhotra P. (2018). Beau's lines. Case Reports, 2018, bcr-2018.
2. Piraccini BM & Iorizzo M. (2007). Drug reactions affecting the nail unit: diagnosis and management. *Dermatologic clinics*, 25(2), 215-221.
3. Daneshgaran G, Dubin, DP, & Gould DJ. (2020). Cutaneous Manifestations of COVID-19: A Systematic Review.

